arXiv:1610.03366v2 [math.DG] 7 Dec 2016

THE KÄHLER-RICCI FLOW ON FANO BUNDLES

XIN FU AND SHIJIN ZHANG

Abstract. We study the behavior of the Kähler-Ricci flow on some Fano bundle which is a trivial bundle on one Zariski open set. We show that if the fiber is $\mathbb{P}^m$ blown up at one point or some weighted projective space blown up at the orbifold point and the initial metric is in a suitable kähler class, then the fibers collapse in finite time and the metrics converge sub-sequentially in Gromov-Hausdorff sense to a metric on the base.

1. Introduction

The Ricci flow, introduced by Hamilton ([18]), has become a powerful tool to study the topology and geometric structures of Riemannian manifolds. In general, the Ricci flow develops finite time singularities. Hamilton's program of Ricci flow with surgeries was carried out by Perelman [23, 24, 25] to prove Thurston's geometrization conjecture. The minimal model theory in birational geometry can be viewed as the complex analogue of Thurston's geometrization conjecture. Later in [4] Cao introduced the Kähler-Ricci flow and use it to prove the existence of Kähler-Einstein metrics on manifolds with negative or vanishing first Chern class ([56, 1]).

There are intensive study of the Kähler-Ricci flow in the past few years. In the general type case, has been studied by Tsuji [55], Tian-Zhang [47], Guo [16], Guo-Song-Weinkove [17], Tian-Zhang [48] and references therein. In the Calabi-Yau fiber case, has been studied by Song-Tian [34], Song-Weinkove [41], Fong-Zhang [13], Gill [15], Tosatti-Weinkove-Yang [52] and references therein. In the Fano case, has been studied by Chen-Tian [5, 6], Chen-Wang [7], Phong-Sturm [26], Phong-Song-Sturm-Weinkove [27, 28, 29], Sesum-Tian [30], Székelyhidi [43], Tian-Zhang [49], Tian-Zhang-Zhang-Zhang [46], Tian-Zhu [50, 51] and references therein.

Moreover, Song-Tian have a nice observation which relate the Kähler-Ricci flow with birational geometry. In [34, 35, 36] they introduced the analytic minimal model program which is parallel to Mori's birational minimal model program. On one hand, Kähler-Ricci flow with surgery can be viewed as the complex analogue of Thurston's three dimensional geometrization conjecture. On the other hand, the surgery is canonical and correspond to the birational surgery in Mori's program such as divisorial contraction or flip, see [2, 19]. In this article, our Fano fiber contraction in metric sense can be viewed as the analogue of Mori fiber space in birational geometry.

In this article we study the behavior of the finite time singularity of the Kähler-Ricci flow. The behavior of the Kähler-Ricci flow with finite time singularity has been studied by Feldman-Ilmanen-Knopf [10], Ilmanen-Knopf[21], Song-Weinkove [38], Song [31, 32], Song-Székelyhidi-Weinkove [33], Tian [44, 45], Zhang [57, 58], Fong [11, 12], Collins-Tosatti [9], La Nave-Tian [22], Song-Yuan [42], Tosatti-Zhang [54] and references therein.

Let (M, g_0) be a compact Kähler manifold of complex dimension $n \geq 2$. We write $\omega_0 = \sqrt{-1}(g_0)_{i\bar{j}} dz^i \wedge d\overline{z^j}$ for the Kähler form associated to g_0. We consider the Kähler-Ricci flow $\omega = \omega(t)$

given by

$$\frac{\partial}{\partial t}\omega = -\mathrm{Ric}(\omega), \qquad \omega(0) = \omega_0, \tag{1.1}$$

where $\mathrm{Ric}(\omega) = -\sqrt{-1}\partial\overline{\partial}\log\det g$, where $g = g(t)$ is the metric associated to ω.

It's well-known that, from Tian-Zhang [47], a maximal smooth solution to (1.1) exists on $[0, T)$ where $T > 0$ is given by

$$T = \sup\{t > 0 | [\omega_0] - 2\pi t c_1(X) > 0\}. \tag{1.2}$$

Song-Székelyhidi-Weinkove [33] studied the behavior of the Kähler-Ricci flow on the projective bundles. One essential point of their proof is that the projective space admits a metric which has positive bisectional curvature. In this article, we want to generalize their result in the sense that we could have more types of Fano fibers other than projective spaces. For example the fiber could be $\mathbb{P}^m$ blown-up at one point or $M_{m,k}$ which is the weighted projective space $Y_{m,k}(1 \leq k < m)$ blown-up at the orbifold point (the definitions of $M_{m,k}$ and $Y_{m,k}$ see section 4). In this paper, we consider the Fano bundle which is trivial on one Zariski open set. More precisely,

Definition 1.1 (Fano bundle)**.** *Let X, Y be compact Kähler manifolds with dimension n, m, respectively, F be a Fano manifold with dimension $n - m$ and a surjective holomorphic map $\pi : X \to Y$. We say X is a Fano bundle over Y with fiber F, if for any $y \in Y$, there exists a Zariski open set $y \in U \subset Y$ and a biholomorphism $\Phi : \pi^{-1}(U) \to U \times F$ such that the diagram*

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\Phi} & U \times F \\ & {\scriptstyle\pi}\searrow \quad \swarrow{\scriptstyle \mathrm{Pr}_1} & \\ & U & \end{array}$$

commutes, where Pr_1 is the projection map onto the first factor. We denote it as (X, Y, π, F).

Remark 1.0.1. *We recall the definition of Fano fibration, was defined by Tosatti-Zhang* [54]*. A compact Kähler manifold is said to admit a Fano fibration if there is a surjective holomorphic map $\pi : X \to Y$ with connected fibers, where Y is a compact normal Kähler space with $0 \leq \dim Y < \dim X$ and such that for every fiber F of π such that $-K_X|_F$ is ample. Hence the Fano bundle is a special Fano fibration.*

Remark 1.0.2. *The simplest example of a Fano bundle is $X = F \times Y$ with a Fano manifold F and any compact Kähler manifold Y. The projective bundle X was considered by Song-Székelyhidi-Weinkove* [33] *is a Fano bundle with $F = \mathbb{P}^{n-m}$.*

Since the fiber F is a Fano manifold, the solution $\omega(t)$ develops a singularity after a finite time. By (1.2), T is finite since $F \cdot c_1(X)^{n-m} > 0$ for every fiber F. Hence we assume that the limiting Kähler class $[\omega_0] - 2\pi T c_1(X)$ satisfies

$$[\omega_0] - 2\pi T c_1(X) = [\pi^* \omega_Y] \tag{1.3}$$

for some Kähler metric ω_Y on Y. It's well-known that the Kähler-Ricci flow equation (1.1) is equivalent to the following complex Monge-Ampère equation

$$\begin{cases} \dfrac{\partial \varphi}{\partial t} = \log \dfrac{(\frac{1}{T}((T-t)\omega_0 + t\pi^*\omega_Y) + \sqrt{-1}\partial\overline{\partial}\varphi)^n}{\Omega} \\ \varphi(0) = 0, \end{cases} \tag{1.4}$$

where Ω is a smooth volume form, $\chi = \sqrt{-1}\partial\overline{\partial}\log\Omega \in -2\pi c_1(X)$, and $\omega(t) = \frac{1}{T}((T-t)\omega_0 + t\pi^*\omega_Y) + \sqrt{-1}\partial\overline{\partial}\varphi > 0$. By Lemma 2.1 below, we know there exists a bound function φ_T on X satisfying $\lim_{t\to T}\varphi(t) = \varphi_T$. We define

$$\omega_T := \pi^*\omega_Y + \sqrt{-1}\partial\overline{\partial}\varphi_T \geq 0. \tag{1.5}$$

Our first main result of this paper is that ω_T is bounded.

Theorem 1.1. *Assume (X, Y, π, F) is a Fano bundle, ω_0 is the Kähler metric on X, ω_Y is a Kähler metric on Y satisfying (1.3) for some $T > 0$, ω_T is defined by (1.5). Then there exists a uniform constant $C > 0$ such that*

$$C^{-1}\pi^*\omega_Y \leq \omega_T \leq C\pi^*\omega_Y. \tag{1.6}$$

We also study the Kähler-Ricci flow on the Fano bundles with the fiber F is $\mathbb{P}^m$ blown up at one point or $M_{m,k}$ which is the weighted projective space $Y_{m,k}$ (the definition see Section 4) blown up at the orbifold point.

The other main result of this paper shows that diameter of manifold X with metric $\omega(t)$ is finite and there exists a sequence of metrics along the Kähler-Ricci flow converge subsequentially to a metric on Y in the Gromov-Hausdorff sense as $t \to T$. It generalizes in some sense the result of Song-Székelyhidi-Weinkove (Theorem 1.1 in [33]).

Theorem 1.2. *Let (X, Y, π, F) be a Fano bundle with F is $\mathbb{P}^m$ blown up at one point ($m \geq 2$) or $F = M_{m,k}(1 \leq k < m)$, ω_Y be a Kähler metric on Y and ω_0 be a Kähler metric on X. Assume $\omega(t)$ is a solution of the Kähler-Ricci flow (1.1) for $t \in [0, T)$ with initial metric ω_0 and $[\omega_0] - 2\pi T c_1(X) = [\pi^*\omega_Y]$, then we have*

1. $\mathrm{diam}(X, \omega(t)) \leq C$ *for some uniform constant $C > 0$;*
2. *There exists a sequence of times $t_i \to T$ and a distance function d_Y on Y (which is uniformly equivalent to the distance induced by ω_Y, such that $(X, \omega(t_i))$ converges to (Y, d_Y) in the Gromov-Hausdorff sense.*

The Theorem 1.2 is a combination of Theorem 3.1 and Theorem 4.1.

When the dimension of X is 2, our method basically can cover most del Pezzo surface. It will be more interesting when the complex structure of the fiber is changing and when the fiber is general Fano variety in higher dimension.

In this paper, the notation 'tr' means that, if $\alpha = \sqrt{-1}\alpha_{i\overline{j}}dz^i \wedge d\overline{z^j}$ is a real $(1,1)$-form then we write

$$\mathrm{tr}_\omega\alpha = g^{i\overline{j}}\alpha_{i\overline{j}} = \frac{n\alpha \wedge \omega^{n-1}}{\omega^n}. \tag{1.7}$$

In this notation, we can write $\Delta f = \mathrm{tr}_\omega(\sqrt{-1}\partial\overline{\partial}f)$.

In section 2, we recall some well known estimates for the Kähler-Ricci flow, and using Song's argument in [32] to establish a estimate of the horizontal level set, as an application, we prove Theorem 1.1. In Section 3, we using the estimate of the horizontal level set established in section 2 and the argument in [33] to prove the case of fiber is $\mathbb{P}^m$ blown up at one point in Theorem 1.2. In Section 4, we recall the definitions of $M_{m,k}$ and $Y_{m,k}$, using He-Sun's theorem [20] that any weighted projective space admits a orbifold Kähler metric with positive bisectional curvature, Song-Weinkove's argument in [40] and the argument of the proof in the case of F is $\mathbb{P}^m$ blown up at one point, we can prove the case of $F = M_{m,k}$ in the Theorem 1.2.

2. The Main Estimates

In this section, we recall some estimates for the Kähler-Ricci flow, establish a estimate for $\omega(t)$ on the horizontal level set and prove the Theorem 1.1.

We define reference $(1,1)-$forms $\hat{\omega}_t$ on X for $t \in [0,T]$ by

$$\hat{\omega}_t = \frac{1}{T}((T-t)\omega_0 + t\pi^*\omega_Y). \tag{2.1}$$

Then $\hat{\omega}_t$ is a Kähler form in the cohomology class $[\omega(t)]$ for $t \in [0,T)$. Let Ω be the unique smooth volume form on X with $\sqrt{-1}\partial\overline{\partial}\log\Omega = \frac{\partial}{\partial t}\hat{\omega}_t =: \chi \in -2\pi c_1(X)$ and $\int_X \Omega = 1$. We also can write $\hat{\omega}_t$ as $\hat{\omega}_t = \omega_0 + t\chi$.

It's well-known that the Kähler-Ricci flow equation (1.1) is equivalent to the following complex Monge-Ampère equation

$$\begin{cases} \dfrac{\partial\varphi}{\partial t} = \log\dfrac{(\hat{\omega}_t + \sqrt{-1}\partial\overline{\partial}\varphi)^n}{\Omega} \\ \varphi(0) = 0 \\ \omega(t) > 0. \end{cases} \tag{2.2}$$

where $\omega(t) = \hat{\omega}_t + \sqrt{-1}\partial\overline{\partial}\varphi$.

In this paper we use C to denote a uniform constant, independent of time but possibly depending on ω_0, n, T, which may differ from line to line. Then the following estimates are well known, see the Lemma 2.1 and Lemma 2.2 in [39].

Lemma 2.1. *For any Kähler manifold (X,ω_0) and Kähler manifold (Y,ω_Y). If there exists a surjective holomorphic map $\pi : X \to Y$, and the smooth solution $\omega(t)$ of the Kähler-Ricci flow (1.1) on X satisfying $\lim_{t\to T}[\omega(t)] = [\pi^*\omega_Y](T < +\infty)$. Then we have*

1. *There exists a uniform constant $C > 0$ such that $||\varphi||_{L^\infty} \leq C$;*
2. *There exists a uniform constant $C > 0$ such that $\dot{\varphi} \leq C$;*
3. *As $t \to T$, $\varphi(t)$ converges pointwise on X to a bounded function φ_T satisfying*

$$\omega_T := \pi^*\omega_Y + \sqrt{-1}\partial\overline{\partial}\varphi_T \geq 0. \tag{2.3}$$

4. *There exists a uniform constant $c > 0$ such that*

$$\omega(t) \geq c\pi^*\omega_Y. \tag{2.4}$$

Next motivated by the argument of Song, see subsection 3.1 in [32], which Song estimated the evolving metrics of the Kähler-Ricci flow in a well-chosen set of directions in the tangent space of each point on X instead of all directions, we estimate the metric $\omega(t)$ on the horizontal level set of the Fano bundle X.

Let (X,Y,π,F) be the Fano bundle (see Definition 1.1). Since for any $x \in X$, let $y = \pi(x)$, there exists a Zariski open set $(y \in)U \subset Y$, such that the diagram

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\Phi} & U \times F \\ & {\scriptstyle\pi}\searrow \quad \swarrow{\scriptstyle \mathrm{Pr}_1} & \\ & U & \end{array}$$

commutes, where Pr_1 is the projection map onto the first factor. Let $f = \mathrm{Pr}_2 \circ \Phi(x)$, $H = \Phi^{-1}(U \times \{f\})$, where Pr_2 is the projection map onto the second factor. Let $D = Y\backslash U$ and s be a

holomorphic section on $[D]$ and let h be a Hermitian metric on $[D]$. Define $|s|^2 = hs\overline{s}$. Then on the horizontal level set H, we have the estimate for $\omega(t)$

Lemma 2.2. *Assume $\omega(t)$ is the solution of the Kähler-Ricci flow (1.1) and $\lim_{t\to T}[\omega(t)] = [\pi^*\omega_Y]$. Fix any point $x \in X$, then there exists $U \subset Y$, let $f(x) = \mathrm{Pr}_2 \circ \Phi(x)$ and $H = \Phi^{-1}(U \times \{f(x)\})$. Then there exist uniform constants $C > 0$ and $\alpha > 0$, such that $\omega(t)|_H \leq \frac{C}{\pi^*(|s|^{2\alpha})}(\pi^*\omega_Y)|_H$.*

Proof. Since for any $x \in \pi^{-1}(U)$, $\pi(x) = y \in U$, and Φ is a biholomorphism from $\pi^{-1}(U)$ to $U \times F$, there exist constants $\alpha > 0$ and $C > 0$ such that $\pi^*\omega_Y|_H \geq \pi^*|s|^\alpha\omega_Y|_U$. On the other hand, for each time $t \in (0, T)$, $\omega(t)$ is equivalent to metric ω_0. Hence if we let

$$u(t, x) = \pi^*(|s|^{2\alpha})\mathrm{tr}_{\pi^*\omega_Y|_H}(\omega(t)|_H)(x),$$

we know $u \to 0$ along $X\backslash\pi^{-1}(U)$ and hence a positive maximum must occur in $\pi^{-1}(U)$ at each fixed time $t \in (0, T)$. We assume the maximum can be obtained at point $x_0 \in X$. Let $y_0 = \pi(x_0) \in Y$. We choose normal coordinate system $(z^i)_{i=1,\cdots,n}$ for $g(t)$ at x_0 and $(w^\alpha)_{\alpha=1,\cdots,m}$ for g_Y at y_0. For any holomorphic vector $\frac{\partial}{\partial w^\alpha}$, there exist holomorphic vector $\frac{\partial}{\partial x^\alpha} \in T_xX$ such that $d\pi_x(\frac{\partial}{\partial x^\alpha}) = \frac{\partial}{\partial w^\alpha}$ for any x in the local normal coordinate chart of x_0. The map π is given locally as $(\pi^1, \cdots, \pi^m)$ for holomorphic functions $\pi^\alpha = \pi^\alpha(z^1, \cdots, z^n)$. We write $\frac{\partial}{\partial x^\alpha}$ as $\frac{\partial}{\partial x^\alpha} = a^i_\alpha\frac{\partial}{\partial z^i}$ for holomorphic functions a^i_α. Then u can be expressed as $u(t, x) = |s|^{2\alpha}(\pi(x))g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{a^j_\beta}g_{i\overline{j}}$. For convenience, we denote $u_1 = g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{a^j_\beta}g_{i\overline{j}}$. Then at point x_0

$$\begin{aligned}
\Delta u_1 &= g^{k\overline{l}}\partial_k\partial_{\overline{l}}(g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{a^j_\beta}g_{i\overline{j}}) \\
&= \sum_{k,l=1}^n g^{k\overline{l}}\partial_k(\partial_{\overline{\delta}}g_Y^{\alpha\overline{\beta}}\overline{\pi^\delta_l}a^i_\alpha\overline{a^j_\beta}g_{i\overline{j}} + g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{\partial_l a^j_\beta}g_{i\overline{j}} + g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{a^j_\beta}\partial_{\overline{l}}g_{i\overline{j}}) \\
&= -\partial_\gamma\partial_{\overline{\delta}}(g_Y)_{\beta\overline{\alpha}}\pi^\gamma_k\overline{\pi^\delta_k}a^i_\alpha\overline{a^i_\beta} + |\partial_k a^i_\alpha|^2 - a^i_\alpha\overline{a^j_\alpha}R_{i\overline{j}} \\
&= (\mathrm{Rm}(g_Y))_{\gamma\overline{\delta}\beta\overline{\alpha}}\pi^\gamma_k\overline{\pi^\delta_k}a^i_\alpha\overline{a^i_\beta} + |\partial_k a^i_\alpha|^2 - a^i_\alpha\overline{a^j_\alpha}R_{i\overline{j}}
\end{aligned}$$

On the other hand,

$$\frac{\partial u_1}{\partial t} = g_Y^{\alpha\overline{\beta}}a^i_\alpha\overline{a^j_\beta}\frac{\partial}{\partial t}g_{i\overline{j}} = -a^i_\alpha\overline{a^j_\alpha}R_{i\overline{j}}.$$

Hence

$$\begin{aligned}
(\frac{\partial}{\partial t} - \Delta)\log u_1 &= \frac{1}{u_1}(-(\mathrm{Rm}(g_Y))_{\gamma\overline{\delta}\beta\overline{\alpha}}\pi^\gamma_k\overline{\pi^\delta_k}a^i_\alpha\overline{a^i_\beta} - |\partial_k a^i_\alpha|^2) + \frac{|\nabla u_1|^2}{u_1^2} \\
&\leq c_Y\mathrm{tr}_\omega\pi^*\omega_Y + \frac{1}{u_1}(\frac{|\nabla u_1|^2}{u_1} - |\partial_k a^i_\alpha|^2),
\end{aligned}$$

where $-c_Y$ is a lower bound for the bisectional curvature of ω_Y on Y. It is easy to get (see [34])

$$\tfrac{|\nabla u_1|^2}{u_1} - |\partial_k a^i_\alpha|^2 \leq 0.$$

Hence we have

$$(\frac{\partial}{\partial t} - \Delta)\log u_1 \leq c_Y\mathrm{tr}_\omega\pi^*\omega_Y. \tag{2.5}$$

Since $\sqrt{-1}\partial\overline{\partial}(\pi^*|s|^2)(x_0) = \sqrt{-1}\partial\overline{\partial}|s|^2(y_0)$, is bounded by some multiple of $\pi^*\omega_Y$. Combine Lemma 2.1, we have

$$(\frac{\partial}{\partial t} - \Delta)\log u \leq C\mathrm{tr}_\omega \pi^*\omega_Y \leq C'.$$

Hence by the maximum principle, we have $u \leq C$. □

Now we prove the Theorem 1.1.

Proof of Theorem 1.1. Lower bound follows from (4) in Lemma 2.1. For any point $y \in Y$, each fiber $\pi^{-1}(y) = F$ is a closed Kähler manifold, and since $\pi^*\omega_Y|_{\pi^{-1}(y)} = 0$, we have

$$\sqrt{-1}\partial\overline{\partial}\varphi_T|_{\pi^{-1}(y)} = \omega_T|_{\varphi_T} \geq 0,$$

since φ_T is bounded, φ_T must be constant on the fiber $\pi^{-1}(y)$. Hence there exists a bounded function ψ_T on Y satisfying

$$\varphi_T = \pi^*\psi_T.$$

Hence

$$\omega_T = \pi^*(\omega_Y + \sqrt{-1}\partial\overline{\partial}\psi_T).$$

Now for any $x \in X$, we may assume that $|s|^2(\pi(x)) = 0$, there exists an open set $\pi(x) \in U \subset Y$, such that Lemma 2.2 holds. Now we consider the open set $U_{1/2} := \{y \in U||s|^2(y) > 1/2\}$. Then by Lemma 2.2, there exists a constant $C > 0$ such that

$$\sqrt{-1}\partial\overline{\partial}\psi_T|_{U_{1/2}} \leq C\omega_Y.$$

Since Y is a compact manifold, there exist a finite open set $\{U^i_{1/2}(1 \leq i \leq N)\}$ (N is a positive integer number) satisfying

$$\cup_{i=1}^N U^i_{1/2} = Y.$$

Hence we obtain that there exists a uniform constant $C > 0$ such that

$$\sqrt{-1}\partial\overline{\partial}\psi_T \leq C\omega_Y.$$

Hence we finish the proof of the theorem. □

3. F Is $\mathbb{P}^m$ Blown Up At One Point

In this section, we consider the case of F is $\mathbb{P}^m$ blown up at one point. One essential point of Song-Székelyhidi-Weinkove's proof [33] is that the projective space admits a metric which has positive holomorphic bisectional curvature. Although $\mathbb{P}^m$ blown up at one point doesn't admit a metric with nonnegative holomorphic bisectional curvature, but we have such metric with nonnegative bisectional curvature on outside of the divisor. Then we need to estimate the locally holomorphic vector field near the divisor under the evolving metrics, by using a idea of Song-Weinkove [39]. We also need Lemma 2.2, estimate of the evolving metrics along the Kähler-Ricci flow which were restricted to a horizontal set. We prove the following

Theorem 3.1. *Let (X, Y, π, F) be a Fano bundle with F is $\mathbb{P}^m$ blown up at one point ($m \geq 2$), ω_Y be a Kähler metric on Y and ω_0 be a Kähler metric on X. Assume $\omega(t)$ is a solution of the Kähler-Ricci flow (1.1) for $t \in [0, T)$ with initial metric ω_0 and $[\omega_0] - 2\pi T c_1(X) = [\pi^*\omega_Y]$, then we have*

(1) $\mathrm{diam}(X, \omega(t)) \leq C$ *for some uniform constant $C > 0$;*

(2) *There exists a sequence of times $t_i \to T$ and a distance function d_Y on Y (which is uniformly equivalent to the distance induced by ω_Y, such that $(X, \omega(t_i))$ converges to (Y, d_Y) in the Gromov-Hausdorff sense.*

3.1. **Key Estimates.** Write $\pi_1 : F \to \mathbb{P}^m$ for the blow-down map, which is an isomorphism from $F\backslash E$ to $\mathbb{P}^m\backslash\{p\}$, where $p \in \mathbb{P}^m$ and $E = \pi_1^{-1}(p)$, which is biholomorphic to $\mathbb{P}^{m-1}$. For convenient, once and for all, a coordinate chart V centered at p, which we identify via coordinates $z^1, \cdots, z^m$ with the unit ball D_1 in $\mathbb{C}^m$,

$$D_1 = \{(z^1, \cdots, z^m) \in \mathbb{C}^m | \sum_{i=1}^m |z^i|^2 < 1\}. \tag{3.1}$$

Denote by g_e the Euclidean metric on D_1. Since g_e and g_{FS} are uniformly equivalent on D_1, it suffices to estimates for g_e on D_1. Write $D_r \subset D_1$ for the ball of radius $0 < r < 1$ with respect to g_e.

We recall the definition of the blow-up construction, following the exposition in [14]. We identify $\pi_1^{-1}(D_1)$ with the submanifold $\tilde{D}_1$ of $D_1 \times \mathbb{P}^{m-1}$ given by

$$\tilde{D}_1 = \{(z, l) \in D_1 \times \mathbb{P}^{m-1} | z^i l^j = z^j l^i\}, \tag{3.2}$$

where $l = [l^1, \cdots, l^m]$ are homogeneous coordinates on $\mathbb{P}^{m-1}$. The map π_1 restricted to $\tilde{D}_1$ is the projection $\pi|_{\tilde{D}_1}(z, l) = z \in D_1$, with the exceptional divisor $E \cong \mathbb{P}^{m-1}$ given by $\pi_1^{-1}(0)$. The map π gives an isomorphism from $\tilde{D}_1\backslash E$ onto the punctured ball $D_1\backslash\{0\}$.

On $\tilde{D}_1$ we have coordinate charts $\tilde{D}_{1i} = \{l^i \neq 0\}$ with local coordinates $\tilde{z}(i)^1, \cdots, \tilde{z}(i)^m$ given by $\tilde{z}(i)^j = l^j/l^i = z^j/z^i$ for $j \neq i$ and $\tilde{z}(i)^i = z^i$. The divisor E is given in $\tilde{D}_{1i}$ by $\{\tilde{z}(i)^i = 0\}$. The line bundle $[E]$ over $\tilde{D}_1$ has transition functions z^i/z^j on $\tilde{D}_{1i} \cap \tilde{D}_{1j}$. We can define a global section s of $[E]$ over F by setting $s(z) = z^i$ on $\tilde{D}_{1i}$ and $s = 1$ on $F\backslash\pi_1^{-1}(D_{1/2})$. The section s_1 vanishes along the exceptional divisor E. We also define a Hermitian metric h_1 on $[E]$ as follows. First let h_2 be the Hermitian metric on $[E]$ over $\tilde{D}_1$ given in $\tilde{D}_{1i}$ by

$$h_2 = \frac{\sum_{j=1}^m |l^j|^2}{|l^i|^2}, \tag{3.3}$$

and let h_3 be the Hermitian metric on $[E]$ over $F\backslash E$ determined by $|s_1|^2_{h_2} = 1$. Now define the Hermitian metric h_1 by $h_1 = \rho_1 h_2 + \rho_2 h_3$, where ρ_1, ρ_2 is a partition of unity for the cover $(\pi_1^{-1}(D_1), F\backslash\pi_1^{-1}(D_{1/2}))$ of F, so that $h_1 = h_2$ on $\pi_1^{-1}(D_{1/2})$. The function $|s_1|^2_{h_1}$ on F is given on $\pi_1^{-1}(D_{1/2})$ by

$$|s_1|^2_{h_1}(x) = \sum_{i=1}^m |z^i|^2 =: r^2, \tag{3.4}$$

for $\pi_1(x) = (z^1, \cdots, z^m)$. On $\pi_1^{-1}(D_{1/2}\backslash\{0\})$, the curvature $R(h_1)$ of h_1 is given by

$$R(h_1) = -\sqrt{-1}\partial\overline{\partial}\log(\sum_{i=1}^m |z^i|^2). \tag{3.5}$$

We have the following lemma (see [14], p.187).

Lemma 3.1. *For sufficiently small $\epsilon_0 > 0$,*

$$\omega_F = \pi_1^*\omega_{FS} - \epsilon_0 R(h_1) \tag{3.6}$$

is a Kähler form on F.

From now on we fix $\epsilon_0 > 0$ as in the Lemma 3.1, with ω_F defined in Lemma 3.1. In $\pi_1^{-1}(D_{1/2}\backslash\{0\})$, which we can identify with $D_{1/2}\backslash\{0\}$, the metric ω_F has the form:

$$\omega_F = \pi_1^*\omega_{FS} + \sqrt{-1}\frac{\epsilon_0}{r^2}\sum_{i,j=1}^{m}(\delta_{ij} - \frac{\overline{z^i}z^j}{r^2})dz^i d\overline{z^j}, \tag{3.7}$$

for r given by (3.4). It is easy to see that, in $D_{1/2}\backslash\{0\}$, $R(h_1) \leq 0$, and the following lemma holds (see [39]).

Lemma 3.2. *There exist positive constants C such*

$$\pi_1^*\omega_{FS} \leq \omega_F \leq C\frac{\pi_1^*\omega_{FS}}{|s_1|^2_{h_1}} \tag{3.8}$$

Since (X, Y, π, F) is a Fano bundle, for any $x \in X$, let $y = \pi(x)$, there exists a Zariski open set $(y \in)U \subset Y$, such that the diagram

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\Phi} & U \times F \\ & {\scriptstyle\pi}\searrow \quad \swarrow{\scriptstyle \mathrm{Pr}_1} & \\ & U & \end{array}$$

commutes, where Pr_1 is the projection map onto the first factor. Let $D = Y\backslash U$ and s be a holomorphic section on $[D]$ and let h be a Hermitian metric on $[D]$. Define $|s|^2 = hs\overline{s}$, for simplicity, we also write $\pi^*|s|^2_h$ as $|s|^2_h$. On $\pi^{-1}(U)$, we denote $\tilde{\omega} = \Phi^*(\mathrm{Pr}_2^*\pi_1^*\omega_{FS} + \mathrm{Pr}_1^*\omega_Y)$, we also write $|s_1|^2_{h_1}$ to represent $\Phi^*\mathrm{Pr}_2^*(|s_1|^2_{h_1})$, where Pr_2 is the projection map onto the second factor. Then we have the following

Lemma 3.3. *There exist uniform constants $C > 0$ and $\alpha > 0$ such that for $\omega = \omega(t)$ a solution of the Kähler-Ricci flow,*

$$\omega(t) \leq \frac{C}{|s|^{2\alpha}_h|s_1|^2_{h_1}}\tilde{\omega}. \tag{3.9}$$

Proof. Fix $0 < \epsilon \leq 1$. By Lemma 3.2, we know

$$\tilde{\omega} \geq C\Phi^*(|s_1|^2_{h_1}(\mathrm{Pr}_2^*\omega_F + \mathrm{Pr}_1^*\omega_Y)).$$

Since $\mathrm{Pr}_2^*\omega_F + \mathrm{Pr}_1^*\omega_Y$ is a fixed Kähler metric on $U \times F$ and Φ is a biholomorphism from $\pi^{-1}(U)$ to $U \times F$, for any fixed time t, there exists a constant $\alpha > 0$ such that

$$\mathrm{tr}_{\tilde{\omega}}\omega \leq \frac{C}{|s|^{\alpha}_h|s_1|^2_{h_1}}.$$

Hence if we set

$$Q_\epsilon = \log(|s|^{2\alpha}_h|s_1|^{2+2\epsilon}_{h_1}\mathrm{tr}_{\tilde{\omega}}\omega). \tag{3.10}$$

For each fixed time $t \in (0, T)$. We know the maximum of Q_ϵ must be obtained at some point $x_0 \in \Phi^{-1}(U \times F\backslash E)$. Now we compute at point (x_0, t)

$$\begin{aligned}(\frac{\partial}{\partial t} - \Delta)Q_\epsilon &= (\frac{\partial}{\partial t} - \Delta)\log \mathrm{tr}_{\tilde{\omega}}\omega + \alpha\mathrm{tr}_\omega R(h) + (1+\epsilon)\mathrm{tr}_\omega R(h_1) \\ &\leq (\frac{\partial}{\partial t} - \Delta)\log \mathrm{tr}_{\tilde{\omega}}\omega + \alpha\mathrm{tr}_\omega R(h).\end{aligned} \tag{3.11}$$

From the argument in the proof of Lemma 2.2, there exists a uniform constant $C > 0$ such that

$$\alpha \mathrm{tr}_\omega R(h) \leq C \mathrm{tr}_\omega \pi^* \omega_Y \leq C'.$$

By a well-known computation (see [56, 1, 4]):

$$\begin{aligned}(\frac{\partial}{\partial t} - \Delta) \log \mathrm{tr}_{\tilde{\omega}} \omega &= \frac{1}{\mathrm{tr}_{\tilde{\omega}} \omega}(-g^{i\overline{j}} \tilde{g}^{k\overline{q}} \tilde{g}^{p\overline{l}} g_{k\overline{l}} \tilde{R}_{i\overline{j}p\overline{q}} - g^{i\overline{j}} \tilde{g}^{k\overline{l}} g^{p\overline{q}} \tilde{\nabla}_i g_{k\overline{q}} \tilde{\nabla}_{\overline{j}} g_{p\overline{l}} + \frac{|\nabla \mathrm{tr}_{\tilde{\omega}} \omega|^2}{\mathrm{tr}_{\tilde{\omega}} \omega}) \\ &\leq -\frac{1}{\mathrm{tr}_{\tilde{\omega}} \omega} g^{i\overline{j}} \tilde{g}^{k\overline{q}} \tilde{g}^{p\overline{l}} g_{k\overline{l}} \tilde{R}_{i\overline{j}p\overline{q}}\end{aligned} \tag{3.12}$$

Denote $\hat{g}$ as the product metric $\mathrm{Pr}_2^* \pi_1^* \omega_{FS} + \mathrm{Pr}_1^* \omega_Y$, then $\tilde{g} = \Phi^* \hat{g}$. We compute with metric $\hat{g}$, since the bisectional curvature of ω_{FS} is positive, we have

$$\hat{R}_{i\overline{j}p\overline{q}} = (\mathrm{Pr}_2^* \pi_1^* R(\omega_{FS}))_{i\overline{j}p\overline{q}} + (\mathrm{Pr}_1^* R(\omega_Y))_{i\overline{j}p\overline{q}} \geq (\mathrm{Pr}_1^* R(\omega_Y))_{i\overline{j}p\overline{q}}. \tag{3.13}$$

Since $\Phi^* \mathrm{Pr}_1^* = \pi^*$, pulling back via the map Φ, we have

$$g^{i\overline{j}} \tilde{g}^{k\overline{q}} \tilde{g}^{p\overline{l}} g_{k\overline{l}} \tilde{R}_{i\overline{j}p\overline{q}} \geq -C(\mathrm{tr}_{\tilde{\omega}} \omega)(\mathrm{tr}_\omega \pi^* \omega_Y)$$

for some uniform constant. Hence we obtain that

$$(\frac{\partial}{\partial t} - \Delta) \log \mathrm{tr}_{\tilde{\omega}} \omega \leq C \mathrm{tr}_\omega \mathrm{Pr}_1^* \omega_Y \leq C'.$$

Hence

$$(\frac{\partial}{\partial t} - \Delta) Q_\epsilon \leq C. \tag{3.14}$$

Then using the maximum principle and letting $\epsilon \to 0$, we obtain the lemma. □

We assume that $|s|_h(y) = 1$ and denote $U_{1/2} = \{\tilde{y} \in U | |s|_h^2(y) > 1/2\}$.
Consider the holomorphic vector field

$$\sum_i^m z^i \frac{\partial}{\partial z^i},$$

defined on the unit ball D_1. This defines via π_1 a holomorphic vector field V on $\pi_1^{-1}(D_1) \subset F$ which vanishes to order 1 along the exceptional divisor E. We can extend V to be a smooth $T^{1,0}$ vector field on the whole of F, and $\mathrm{Pr}_2^*(V)$ to be a smooth $T^{1,0}$ vector field on $U_{1/2} \times F$, then pull back by Φ and then extend it to a vector $\tilde{V}$ on whole of X which vanish on $X \backslash \pi^{-1}(U_{1/2})$. We then have the following lemma.

Lemma 3.4. *For $\omega = \omega(t)$ a solution of the Kähler-Ricci flow, we have the estimate*

$$|\tilde{V}|_\omega^2 \leq C |s_1|_{h_1}, \tag{3.15}$$

for a uniform constant C. Locally, in $D_{1/2} \backslash \{0\}$ we have

$$|W|_g^2 \leq \frac{C}{r}, \tag{3.16}$$

for

$$W = \sum_{i=1}^m (\frac{x^i}{r} \frac{\partial}{\partial x^i} + \frac{y^i}{r} \frac{\partial}{\partial y^i})$$

the unit length radial vector field with respect to g_e, where $z^i = x^i + \sqrt{-1} y^i$.

In the statement and proof of the lemma, we are identifying $\pi_1^{-1}(D_{1/2}\backslash\{0\})$ with $D_{1/2}\backslash\{0\}$ via the map π_1, writing g for the Kähler metric $(\pi_1^{-1})^*(\omega|_F)$.

Proof. In this proof, we denote ω_U for $\Phi^*(\mathrm{Pr}_1^*\omega_Y + \mathrm{Pr}_2^*\omega_F)$ on $\pi^{-1}(U)$ and a Hermitian metric $\tilde{\omega} = \rho_1\omega_0 + \rho_2\omega_U$, where ρ_1, ρ_2 is a partition of unity for the cover $(X\backslash\pi^{-1}(U_{1/2}), \pi^{-1}(U))$, so that $\tilde{\omega} = \omega_U$ on $\pi^{-1}(U_{1/2})$ and which is equivalent to metric ω_0.

From the Lemma 3.1 we have, in $D_{1/2}$,

$$|\tilde{V}|^2_{\tilde{\omega}} = |V|^2_{\omega_F} = |V|^2_{\pi^*\omega_{FS}}. \tag{3.17}$$

It follows that $|\tilde{V}|^2_{\omega_0}$ is uniformly equivalent to $|s_1|^2_{h_1} = r^2$ in $D_{1/2}$.

For any fixed point (x,t). We compute in a normal coordinate system for g at (x,t), we have

$$(\frac{\partial}{\partial t} - \Delta)\log|\tilde{V}|^2_{\omega} = \frac{1}{|\tilde{V}|^2_{\omega}}(-g^{i\overline{j}}g_{k\overline{l}}(\partial_i\tilde{V}^k)(\overline{\partial_j\tilde{V}^l}) + \frac{|\nabla|\tilde{V}|^2_{\omega}|^2_{\omega}}{|\tilde{V}|^2_{\omega}}) \leq 0. \tag{3.18}$$

Where we use the Cauchy-Schwarz inequality to get the above inequality (the detail, see the proof of Lemma 2.6 in [39]).

Then using the maximum principle, we obtain that there exists a positive constant C such that

$$|\tilde{V}|^2_{\omega} \leq C|s_1|^2_{h_1}. \tag{3.19}$$

Now define a Hermitian metric $\tilde{\omega}_F$ on $\mathbb{P}^m$ by

$$\tilde{\omega}_F = \omega_e \quad \text{on} \quad D_1,$$

and extending in an arbitrary way to be a smooth Hermitian metric on F. For small $\epsilon > 0$, we consider the quantity

$$Q_\epsilon = \log(|\tilde{V}|^{2+2\epsilon}_{\omega}|s|^{2\alpha+2\epsilon}_h \mathrm{tr}_{\Phi^*(\mathrm{Pr}_2^*\pi_1^*\tilde{\omega}_F + \mathrm{Pr}_1^*\omega_Y)}\omega) - At \tag{3.20}$$

where α is the constant in Lemma 3.3 and A is a constant to be determined. Since $\tilde{\omega}_F$ is uniformly equivalent to ω_{FS}, we see that for fixed t, using Lemma 3.3 and (3.19),

$$(|V|^{2+2\epsilon}_{\omega}|s|^{2\alpha+2\epsilon}_h \mathrm{tr}_{\Phi^*(\mathrm{Pr}_2^*\pi_1^*\tilde{\omega}_F + \mathrm{Pr}_1^*\omega_Y)}\omega)(x)$$

tends to zero as x tends to $X\backslash\pi^{-1}(U_{1/2}) \cup \Phi^{-1}(U_{1/2}\times E)$ and thus $Q_\epsilon(x)$ tends to negative infinity. We now applying the maximum principle to Q_ϵ. Since Q_ϵ is uniformly bounded from above on $\Phi^{-1}(U_{1/2}\times F)\backslash\Phi^{-1}(U_{1/2}\times D_{1/2}\backslash\{0\})$, we only need to rule out the case when Q_ϵ attains its maximum at a point in $\Phi^{-1}(U_{1/2}\times D_{1/2}\backslash\{0\})$. Assume that at some point $(x_0,t_0) \in \Phi^{-1}(U_{1/2}\times D_{1/2}\backslash\{0\})\times(0,T)$, we have $\sup_{\Phi^{-1}(U\times F\backslash E)\times[0,t_0]} Q_\epsilon = Q_\epsilon(x_0,t_0)$.

As in the proof of Lemma 3.3, we have

$$(\frac{\partial}{\partial t} - \Delta)\log|s|^{2\alpha+2\epsilon}_h = (\alpha+\epsilon)\mathrm{tr}_\omega R(h) \leq C\mathrm{tr}_\omega\pi^*\omega_Y. \tag{3.21}$$

By (3.12), pulling back by the biholomorphic map Φ, we have in $\Phi^{-1}(U_{1/2}\times D_{1/2}\backslash\{0\})$,

$$(\frac{\partial}{\partial t} - \Delta)\log(\mathrm{tr}_{\Phi^*(\mathrm{Pr}_2^*\pi_1^*\tilde{\omega}_F + \mathrm{Pr}_1^*\omega_Y)}\omega) \leq C\mathrm{tr}_\omega\Phi^*\mathrm{Pr}_1^*\omega_Y. \tag{3.22}$$

Hence

$$(\frac{\partial}{\partial t} - \Delta)\log|s|^{2\alpha+2\epsilon}_h(\mathrm{tr}_{\Phi^*(\mathrm{Pr}_2^*\pi_1^*\tilde{\omega}_F + \mathrm{Pr}_1^*\omega_Y)}\omega) \leq C\mathrm{tr}_\omega\pi^*\omega_Y \leq C'. \tag{3.23}$$

Take $A = C' + 1$. By (3.18), at (x_0, t_0), we obtain

$$0 \leq (\frac{\partial}{\partial t} - \Delta)Q_\epsilon \leq -1, \tag{3.24}$$

a contradiction. Thus Q_ϵ is uniformly bounded from above. Letting ϵ tend to zero, since $|s|_h^2 > 1/2$ on $\pi^{-1}(U_{1/2})$, we obtain

$$(\mathrm{tr}_{\Phi^*(\mathrm{Pr}_2^*\pi_1^*\omega_{FS}+\mathrm{Pr}_1^*\omega_Y)}\omega)|\tilde{V}|_\omega^2 \leq C, \tag{3.25}$$

for some uniform constant C. By Lemma 3.2, we have

$$(\mathrm{tr}_{\omega_0}\omega)|\tilde{V}|_\omega^2 \leq C,$$

and since $|\tilde{V}|_\omega^2 \leq (\mathrm{tr}_{\omega_0}\omega)|\tilde{V}|_{\omega_0}^2$ this gives

$$|\tilde{V}|_\omega^4 \leq C|\tilde{V}|_{\omega_0}^2, \tag{3.26}$$

then the lemma follows from the fact that $|\tilde{V}|_{\omega_0}^2$ is uniformly equivalent to $|s_1|_{h_1}^2$ in $D_{1/2}$. □

Next we estimate on the lengths of spherical and radial paths in the punctured ball $D_{1/2}\backslash\{0\}$, which again we identify with its preimage in each fiber under π_1.

Lemma 3.5. *We have*

1. *For any $y \in Y$ and for $0 < r < 1/2$, the diameter of the $2m-1$ sphere S_r of radius r in D_1 centered at the origin with the metric induced from $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above, independent of r and y.*
2. *For any $z \in D_{1/2}\backslash\{0\}$, the length of a radial path $\gamma(\lambda) = \lambda z$ for $\lambda \in (0, 1]$ with respect to $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above by a uniform constant multiple of $|z|^{1/2}$.*

Hence the diameter of $D_{1/2}\backslash\{0\}$ with respect to $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above and

$$\mathrm{diam}(\pi^{-1}(y), \omega|_{\pi^{-1}(y)}) \leq C.$$

Proof. For any $y \in Y$, we can choose $|s|_h^2(y) = 1$. Then using Lemma 3.3, consider the metric $\omega|_{\pi^{-1}(y)}$, we have

$$\omega|_{\pi^{-1}(y)} \leq \frac{C}{|s_1|_{h_1}^2}\pi_1^*\omega_{FS}. \tag{3.27}$$

Then using the same argument in the proof of Lemma 2.7 in [39], we obtain the lemma. □

Now we can prove the (1) of Theorem 3.1.

Proof of (1) in Theorem 3.1. For any $p, q \in X$. Denote $p_1 = \pi(p), q_1 = \pi(q)$. Then there exist two open subset $U_1, U_2 \subset Y$, such that $p_1 \in U_1, p_2 \in U_2$ and there exist holomorphic maps Φ_1, Φ_2 such that $\Phi_i : \pi^{-1}(U_i) \to U_i \times F (i = 1, 2)$ are the biholomorphic map. Denote $p_2 = \mathrm{Pr}_2\Phi_1(p), q_2 = \mathrm{Pr}_2\Phi_2(q)$. Since Y is compact, we may assume $U_1 \cap U_2 \neq \varnothing$. We assume $\tilde{p}_1 \in U_1 \cap U_2$, denote $\tilde{p} = \Phi_1^{-1}((\tilde{p}_1, p_2))$ and $\tilde{q} = \Phi_2^{-1}((\tilde{p}_1, q_2))$, by Lemma 2.2 we know $d_{\omega(t)}(p, \tilde{p}) \leq C$ and $d_{\omega(t)}(q, \tilde{q}) \leq C$.

Since $\tilde{p}, \tilde{q} \in \pi^{-1}(\tilde{p}_1)$, then by Lemma 3.5, we have $d_{\omega(t)}(\tilde{p}, \tilde{q}) \leq C$. Using the triangle inequality we finish the proof of (1) in Theorem 1.2. □

3.2. **Diameter of fiber tends to zero.** In this subsection, we prove that the diameter of fiber with $\omega(t)$ tends to zero as $t \to T$. Let $d_\omega = d_{\omega(t)}$ be the distance function on X associated to the evolving Kähler metric ω. Using the same argument in the proof of Lemma 3.2 and Lemma 3.3 in [39], we prove the following lemmas.

Lemma 3.6. *Fix a point $y_0 \in Y$. There exists a uniform constant C (independent of y_0) such that for any $p, q \in E$, and any $t \in [0, T)$,*

$$d_\omega(\Phi^{-1}(y_0, p), \Phi^{-1}(y_0, q)) \leq C(T-t)^{1/3}. \tag{3.28}$$

Proof. We can assume that $|s|_h^2(y_0) = 1$. We replace E by $\Phi^{-1}(\{y_0\} \times E)$ in the proof of Lemma 3.2 in [39], using Lemma 3.3 and using Lemma 3.5. See the argument of the proof of Lemma 3.2 in [39]. □

Combine Lemma 3.5 and Lemma 3.6, we have

Lemma 3.7. *Fix a point $y_0 \in Y$. There exists a uniform constant C (independent of y_0) such that for any $0 < \delta_0 < 1/2$ and for anyt $\in [0, T)$*

$$\mathrm{diam}_{\omega(t)}(\Phi^{-1}(\{y_0\} \times \pi_1^{-1}(D_{\delta_0}))) < C(|\delta_0|^{1/2} + (T-t)^{1/3}). \tag{3.29}$$

Proof. We also assume that $|s|_h^2(y_0) = 1$. For any $p, q \in \pi_1^{-1}(D_{\delta_0}))$. Since Lemma 3.6, we only consider $p \in \pi_1^{-1}(D_{\delta_0} \backslash \{0\}))$ and $q \in E$. By Lemma 3.5 (2), we know the length of a radial path $\gamma(\lambda) = \lambda p$ with respect to ω is uniformly bounded from above by $C|p|^{1/2} \leq C\delta_0^{1/2}$. Since $\gamma(\lambda)$ tends to a point $p_0 \in E$ as $\lambda \to 0^+$, we know $d_{\omega|_{\pi^{-1}(y_0)}}(p, p_0) \leq C\delta_0^{1/2}$. By Lemma 3.6, $d_\omega(\Phi^{-1}((y_0, p_0)), \Phi^{-1}((y_0, q))) \leq C(T-t)^{1/3}$. Hence we have

$$d_\omega(\Phi^{-1}((y_0, p)), \Phi^{-1}((y_0, q))) \leq C(\delta_0^{1/2} + (T-t)^{1/3}).$$

□

Lemma 3.8. *Fix a point $y_0 \in Y$. There exists a uniform constant C (independent of y_0) such that for any $p, q \in \pi^{-1}(y_0)$, and any $t \in [0, T)$,*

$$d_\omega(p, q) \leq C(T-t)^{1/15}. \tag{3.30}$$

Proof. We also assume that $|s|_h^2(y_0) = 1$. For each fixed t satisfying $T - t < 2^{-15}$, i.e., $2(T-t)^{2/15} < 1/2$. Take $\delta_0 = (T-t)^{2/15}$, by Lemma 3.7, we have

$$\mathrm{diam}_{\omega(t)}(\Phi^{-1}(\{y_0\} \times \pi_1^{-1}(D_{2\delta_0}))) < C((T-t)^{1/15} + (T-t)^{1/3}) \leq C'(T-t)^{1/15}.$$

We denote $p' = \mathrm{Pr}_2 \circ \Phi(p), q' = \mathrm{Pr}_2 \circ \Phi(p)$. Hence we only consider the case of $p', q' \in F \backslash \pi_1^{-1}(D_{\delta_0})$.

Since $\pi^{-1}(y_0)$ is biholomorphic to F, which is $\mathbb{P}^m$ blown up at one point, there exists a curve $\gamma \cong \mathbb{P}^1$, such that $p, q \in \gamma \cap (\pi^{-1}(y_0) \backslash \Phi^{-1}(\{y_0\} \times \pi_1^{-1}(D_{\delta_0})))$. We may assume that p, q lie in a fixed coordinate chart U whose image under the holomorphic coordinate $z = x + \sqrt{-1}y$ is a ball of radius 2 in $\mathbb{C} = \mathbb{R}^2$ with respect to the Euclidean metric ω_e. In this coordinate, by Lemma 3.3, we know

$$\omega(t)|_{\pi^{-1}(y_0)} \leq \frac{C}{(\delta_0)^2}(\pi_1)^* \omega_{FS} \leq \frac{C'}{(\delta_0)^2}\omega_e.$$

Since closed curve $\gamma \subset F$,

$$\int_\gamma \omega(t) = \int_\gamma [\frac{1}{T}((T-t)\omega_0 + t\pi^*\omega_Y)] = \frac{T-t}{T}\int_\gamma \omega_0 \leq C(T-t). \tag{3.31}$$

Write $\sigma = (T-t)^{1/3} > 0$, which we may assume is sufficiently small.

Moreover, we may assume that p is represented by the origin in $\mathbb{C} = \mathbb{R}^2$, that q is represented by the point $(x_0, 0)$ with $0 < x_0 < 1$, and that the rectangle

$$\mathcal{R} = \{(x,y) \in \mathbb{R}^2 | 0 \leq x \leq x_0, -\sigma \leq y \leq \sigma\} \subset \mathbb{R}^2 = \mathbb{C}$$

is contained in the image of U. Now in $\mathcal{R}$, the fixed metric $\hat{g}_0$ induced from the metric g_0 on X is uniformly equivalent to the Euclidean metric. Thus from (3.31),

$$\int_{-\sigma}^{\sigma} (\int_0^{x_0} (\mathrm{tr}_{\hat{g}_0} g) dx) dy = \int_{\mathcal{R}} (\mathrm{tr}_{\hat{g}_0} g) dx dy \leq C(T-t). \tag{3.32}$$

Hence there exists $y' \in (-\sigma, \sigma)$ such that

$$\int_0^{x_0} (\mathrm{tr}_{\hat{g}_0} g)(x, y') dx \leq \frac{C}{\sigma}(T-t) = C(T-t)^{2/3}. \tag{3.33}$$

Now let p'' and q'' be the points represented by coordinates $(0, y')$ and (x_0, y'). Then, considering the horizontal path $s \mapsto (s, y')$ between p'' and q'', we have

$$\begin{aligned} d_\omega(p'', q'') &\leq \int_0^{x_0} (\sqrt{g(\partial_x, \partial_x)})(x, y') dx \\ &= \int_0^{x_0} (\sqrt{\mathrm{tr}_{\hat{g}_0} g} \sqrt{\hat{g}_0(\partial_x, \partial_x)})(x, y') dx \\ &\leq (\int_0^{x_0} (\mathrm{tr}_{\hat{g}_0} g)(x, y') dx)^{1/2} (\int_0^{x_0} (\hat{g}_0(\partial_x, \partial_x))(x, y') dx)^{1/2} \\ &\leq C(T-t)^{1/3}. \end{aligned} \tag{3.34}$$

and

$$d_\omega(p, p'') \leq d_{\omega|_{\pi^{-1}(y_0)}}(p, p'') \leq \frac{C}{(\delta_0)^2}\sigma = C(T-t)^{1/15}. \tag{3.35}$$

Using the same argument we can prove

$$d_\omega(q, q'') \leq C(T-t)^{1/15}. \tag{3.36}$$

Hence by triangle inequality

$$d_{\omega(t)}(p, q) \leq C(T-t)^{1/15}. \tag{3.37}$$

□

3.3. **Gromov-Hausdorff Convergence.** In this subsection, we prove that there exists a sequence of metrics along the Kähler-Ricci flow converges sub-sequentially to a metric on Y in the Gromov-Hausdorff sense as $t \to T$.

Lemma 3.9. *Write $d_t : X \times X \to \mathbb{R}$ for the distance function induced by the metric $\omega(t)$. There exists a sequence of times $t_i \to T$, such that the functions d_{t_i} converge uniformly to a function $d_\infty : X \times X \to \mathbb{R}$.*

Moreover, if, for $p, q \in Y$, we let $d_{Y,\infty}(p,q) = d_\infty(\tilde{p}, \tilde{q})$, where $\tilde{p} \in \pi^{-1}(p)$ and $\tilde{q} \in \pi^{-1}(q)$, then $d_{Y,\infty}$ defines a distance function on Y, which is uniformly equivalent to that induced by ω_Y.

Proof. First note that the functions $d_t : X \times X \to \mathbb{R}$ are uniformly bounded. Indeed by (1) in Theorem 1.2, we have a constant $C > 0$ such that $d_t(x, y) < C$ for any $t < T$ and $x, y \in X$. Next, we prove that the functions $d_t : X \times X \to \mathbb{R}$ are equicontinuous with respect to the metric on $X \times X$ induced by d_0.

For any $x, x', y, y' \in X$, we have

$$\begin{aligned} |d_t(x, y) - d_t(x', y')| &\leq |d_t(x, y) - d_t(x, y')| + |d_t(x, y') - d_t(x', y')| \\ &\leq d_t(y, y') + d_t(x, x'). \end{aligned} \tag{3.38}$$

Define $|s|_h^2 = hs\overline{s}$, and we assume $|s|_h(y) = 1$.

Since Y is compact, there exist finite Zariski open sets $U^1, \cdots, U^N$ and biholomorphic map $\Phi_1, \cdots, \Phi_N$ such that the diagram

$$\begin{array}{ccc} \pi^{-1}(U^i) & \xrightarrow{\Phi_i} & U^i \times F \\ & {\scriptstyle \pi} \searrow \quad \swarrow {\scriptstyle \mathrm{Pr}_1} & \\ & U^i & \end{array}$$

commutes, where $D_i = Y \backslash U^i$, and $\cup_{i=1}^N \pi^{-1}(U^i_{1/3}) = X$. Let s_i be the holomorphic sections on $[D_i]$ and let h_i be the Hermitian metrics on $[D_i]$. Here $U^i_r = \{\tilde{y} \in U^i || s_i|_{h_i} > r\}$ for $0 < r \leq 1$.

Claim 3.1.1. *There exists a uniform constant $\delta > 0$ such that if $x \in \pi^{-1}(U^{i_0}_{1/3})$ for some $i_0 \in \{1, \cdots, N\}$ and $d_0(x, x') < \delta$, then $x' \in \pi^{-1}(U^{i_0}_{2/3})$.*

Proof of Claim. Denote A^i_r be the boundary of $\pi^{-1}(U^i_r)$ and denote $d_0(A^i_{1/3}, A^i_{2/3}) = \delta_i > 0$. Let ρ^i_1, ρ^i_2 be the partition of unity for the cover $(X \backslash \pi^{-1}(U^i_{2/3}), \pi^{-1}(U^i)$. Then $\omega^i = \rho^i_1 \omega_0 + \rho^i_2 \Phi_i^*(\mathrm{Pr}_1^* \omega_Y + \mathrm{Pr}_2^* \omega_F)$ are the Hermitian metrics on X, which are equivalent to metric ω_0, and $\omega^i = \Phi_i^*(\mathrm{Pr}_1^* \omega_Y + \mathrm{Pr}_2^* \omega_F)$ on $\pi^{-1}(U^i_{2/3})$. Hence there exists a uniform constant $C > 0$ such that

$$C^{-1}\omega_0 \leq \omega^i \leq C\omega_0. \tag{3.39}$$

We take $\delta = C^{-2} \min\{\delta_1, \cdots, \delta_N\}$. Now we can prove $x' \in \pi^{-1}(U^{i_0}_{2/3})$. If not, we know

$$d_{\omega^{i_0}}(x, x') \geq d_{\omega^{i_0}}(A^{i_0}_{1/3}, A^{i_0}_{2/3}) \geq C^{-1} d_0(A^{i_0}_{1/3}, A^{i_0}_{2/3}) = C^{-1}\delta_{i_0}. \tag{3.40}$$

On the other hand,

$$d_{\omega^{i_0}}(x, x') \leq C d_0(x, x') < C\delta. \tag{3.41}$$

It is a contradiction. Hence $x' \in \pi^{-1}(U^{i_0}_{2/3})$. We finish the proof of the claim. □

Now if $x, x' \in X$ satisfying $d_0(x, x') < \delta$, by the above claim we have $x, x' \in U^{i_0}_{2/3}$ for some $i_0 \in \{1, \cdots, N\}$. Now we choose $q_0 \in F$ such that $|s|_h^2(q_0) = 1$. Then by the Lemma 3.3, we have

$$d_t(\Phi_{i_0}^{-1}(\pi(x), q_0), \Phi_{i_0}^{-1}(\pi(x'), q_0)) \leq C d_{\omega_Y}(\pi(x), \pi(x')) \leq C' d_0(x, x'). \tag{3.42}$$

By Lemma 3.8

$$\begin{aligned} d_t(x, x') &\leq d_t(x, \Phi_{i_0}^{-1}(\pi(x), q_0)) + d_t(\Phi_{i_0}^{-1}(\pi(x'), q_0), x') + d_t(\Phi_{i_0}^{-1}(\pi(x), q_0), \Phi_{i_0}^{-1}(\pi(x'), q_0)) \\ &\leq C[(T - t)^{1/15} + d_0(x, x')]. \end{aligned} \tag{3.43}$$

Now we prove the following lemma.

Lemma 3.10. *With the assumption of* (3.43)*, there exists a sequence of times $t_i \to T$, such that the functions d_{t_i} converges uniformly to a continuous function d_∞.*

Proof of Lemma 3.10. We denote $M = X \times X$, is a compact manifold. The first thing to recall is that any compact metric space has a countable dense subset. This follows directly from the definition of compactness. Namely, given any $k \in \mathbb{N}$, cover M by all the balls of radius $\frac{1}{k}$ (centred at all the points of M.) By compactness of M this has a finite subcover, let $Q_k \subset M$ be the set of centers of such a finite subcover. Then every point of M is in one of the balls, so it is distant at most $\frac{1}{k}$ from (at least) one of the points in Q_k. The union, $Q = \cup_k Q_k$, of these finite sets is (at most) countable and is clearly dense in M, i.e., any point in M is the limit of a sequence in Q.

Let $\{d_{t_n}\}$ be a sequence of d_t ($t_n \to T$ as $n \to \infty$). Take a point $q_1 \in Q$, then $\{d_{t_n}(q_1)\}$ is a bounded sequence in $\mathbb{R}$. So, by Heine-Borel Theorem, we may extract a subsequence of d_{t_n} so that $\{d_{t_{n_{1,j}}}(q_1)\}$ converges in $\mathbb{R}$. Since Q is countable we can construct successive subsequences, $d_{t_{n_{k,j}}}$ of the preceding subsequence $d_{t_{n_{k-1,j}}}$, so that the kth subsequence converges at the first kth point $\{q_1, \cdots, q_k\}$ of Q. Now, the diagonal sequence $d_{t_{n_i}} = d_{t_{n_{i,i}}}$ is a subsequence of d_{t_n}. So along this subsequence $d_{t_{n_i}}(q)$ converges for each point in Q. It is a subsequence of the original sequence $\{d_{t_n}\}$, we just denote it as $\{d_{t_n}\}$ and we want to show that it converges uniformly; it suffices to show that it is uniformly Cauchy.

For any given $\epsilon > 0$. By the assumption of (3.43), we can choose $T_\epsilon \in [0, T)$ such that for any $t \in [T_\epsilon, T)$ we have $C(T-t)^{1/15} \leq \epsilon/6$ and choose $\delta = \epsilon/6C > 0$ so that $|d_{t_n}(x) - d_{t_n}(y)| < \epsilon/3$ whenever $d_0(x, y) < \delta$ and $t_n \in [T_\epsilon, T)$. Next choose $k > 1/\delta$. Since there are only finitely many points in Q_k we may choose N so large that for any $n > N$ we have $t_n \in [T_\epsilon, T)$, then $|d_{t_n}(q) - d_{t_m}(q)| < \epsilon/3$ if $q \in Q_k$ and $n, m > N$. Then for a general point $x \in M$ there exists $q \in Q_k$ with $d_0(x, q) < 1/k < \delta$, so

$$|d_{t_n}(x) - d_{t_m}(x)| \leq |d_{t_n}(x) - d_{t_n}(q)| + |d_{t_n}(q) - d_{t_m}(q)| + |d_{t_m}(q) - d_{t_m}(x)| < \epsilon \tag{3.44}$$

whenever $n, m > N$. Thus the sequence is uniformly Cauchy, hence uniformly convergent. Hence the function d_∞ is continuous. We finish the proof of Lemma 3.10. □

It follows that d_∞ is nonnegative, symmetric and satisfies the triangle inequality.

Let $d_Y : Y \times Y \to \mathbb{R}$ be the distance function on Y induced by the metric ω_Y. From Lemma 2.1, we have a constant $c > 0$ such that $d_t(x, y) \geq \sqrt{c} d_Y(\pi(x), \pi(y))$. It follows that the limit d_∞ satisfies

$$d_\infty \geq \sqrt{c} d_Y(\pi(x), \pi(y)). \tag{3.45}$$

Now we want to prove the upper bound. When $\pi(x), \pi(y) \in U^{i_0}_{1/3}$ for some $i_0 \in \{1, \cdots, N\}$, by the inequality (3.43) and Lemma 3.5, we have

$$d_t(x, y) \leq C(T-t)^{1/15} + C d_Y(\pi(x), \pi(y)). \tag{3.46}$$

This implies that

$$d_\infty(x, y) \leq C d_Y(\pi(x), \pi(y)). \tag{3.47}$$

Now we consider the general case. Assume $\pi(x) \in U^1_{1/3}$ and $\pi(y) \in U^{i_0}_{1/3}$ for some $i_0 \neq 1$. Since $\cup_{i=1}^N U^i_{1/3} = Y$, we know for any minimal geodesic $\gamma(t)$ connecting $\pi(x)$ and $\pi(y)$ with metric ω_Y, there exist a finite points $y_0 = \pi(x), y_1 = \gamma(t_1), y_2 = \gamma(t_2), \cdots, y_L = \gamma(t_L), y_{L+1} = \pi(y) \in Y$ such

that y_i and y_{i+1} are in the same $U_{1/3}^{j_0}$ for some $j_0 \in \{1, 2, \cdots, N\}$. We choose $x_0 = x, x_{L+1} = y$ and any $x_1, \cdots, x_L \in X$ satisfying $\pi(x_i) = y_i$. Hence by

$$\begin{aligned} d_Y(\pi(x), \pi(y)) &= \sum_{i=0}^{L} d_Y(y_i, y_{i+1}) \\ &\geq C^{-1} \sum_{i=0}^{L} d_\infty(x_i, x_{i+1}) \\ &\geq C^{-1} d_\infty(x, y). \end{aligned} \tag{3.48}$$

For $p, q \in Y$, we now define $d_{Y,\infty}(p, q) = d_\infty(\tilde{p}, \tilde{q})$, where $\tilde{p} \in \pi^{-1}(p), \tilde{q} \in \pi^{-1}(q)$. This is independent of the choice of lifts $\tilde{p}$ and $\tilde{q}$ since if say $\tilde{p}'$ is a different lift of p, then by (3.48) and by the triangle inequality, we have

$$d_\infty(\tilde{p}', \tilde{q}) \leq d_\infty(\tilde{p}, \tilde{q}) + d_\infty(\tilde{p}', \tilde{p}) = d_\infty(\tilde{p}, \tilde{q}) \tag{3.49}$$

and by switching $\tilde{p}$ and $\tilde{p}'$ we get the reverse inequality. Moreover, it follows from (3.47) and (3.48) that $d_{Y,\infty}$ is uniformly equivalent to d_Y. Hence we finish the proof of Lemma 3.9. □

Theorem 3.2. *In the notation of Lemma 3.9 we have $(X, d_{t_i}) \to (Y, d_{Y,\infty})$ in the Gromov-Hausdorff sense, where we recall that d_{t_i} is the distance function induced by the metric $\omega(t_i)$.*

Proof. Using the same argument in the proof of Theorem 3.1 in [33]. Hence finish the proof of (2) in Theorem 3.1. □

4. F Is Some Weighted Projective Space Blown Up At The Orbifold Point

In this section, we will consider the case of the fiber F is the family of m-folds $M_{m,k} (1 \leq k < m)$, was introduced by Calabi [3], which as generalization of the Hirzebruch surfaces. $M_{n,k}$ is a compactification of the blow up of a $\mathbb{Z}_k$-orbifold point of the orbifold $Y_{m,k}$, is a $\mathbb{P}^1$-bundle over $\mathbb{P}^{m-1}$. The detail of the construction of $M_{m,k}$ and $Y_{m,k}$, please see [40].

First, we recall the definitions of $M_{m,k}$ and $Y_{m,k}$.

We define $M_{m,k}$ to be the $\mathbb{P}^1$-bundle

$$M_{m,k} = \mathbb{P}(\mathcal{O}(-k) \oplus \mathcal{O}) \tag{4.1}$$

over $\mathbb{P}^{n-1}$. We will assume in this paper that $m \geq 2$ and $1 \leq k < m$ (the latter implies that $M_{m,k}$ is a Fano manifold). Denote by E_0 and E_∞ the divisors in $M_{m,k}$ corresponding to sections of $\mathcal{O}(-k) \oplus \mathcal{O}$ with zero $\mathcal{O}(-k)$ and $\mathcal{O}$ component, respectively (the detail see Section 9 in [40]). E_0 is an exceptional divisor with normal bundle $\mathcal{O}(-k)$ of the type discussed above. The complex manifold $M_{m,k}$ can be described as

$$\begin{aligned} M_{m,k} = &\{([Z_1, \cdots, Z_m], (\sigma, \mu) \in \mathbb{P}^{m-1} \times ((\mathbb{C}^m \times \mathbb{C}) \backslash \{(0,0)\}) | \sigma \text{ is in} \\ &\text{the line } \lambda \mapsto (\lambda(Z_1)^k, \cdots, \lambda(Z_m)^k)\} / \sim, \end{aligned} \tag{4.2}$$

where

$$([Z_1, \cdots, Z_m], (\sigma, \mu)) \sim ([Z_1, \cdots, Z_m], (\sigma', \mu')) \tag{4.3}$$

if there exists $a \in \mathbb{C}^*$ such that $(\sigma, \mu) = (a\sigma', a\mu')$. Then E_0 and E_∞ are the divisors $\{\sigma = 0\}$ and $\{\mu = 0\}$, respectively.

The orbifold $Y_{m,k}$ is the weighted projective space

$$Y_{m,k} = \{(Z_0, \cdots, Z_m) \in \mathbb{C}^{n+1}\}/\sim . \tag{4.4}$$

where $(Z_0', \cdots, Z_m') \sim (Z_0, \cdots, Z_m)$ if there exists $\lambda \in \mathbb{C}^*$ such that

$$(Z_0', \cdots, Z_m') = (\lambda^k Z_0, \lambda Z_1, \cdots, \lambda Z_m). \tag{4.5}$$

We write elements of $Y_{m,k}$ as $[Z_0, \cdots, Z_m]$. Then $Y_{m,k}$ has a single $\mathbb{Z}_k$-orbifold point at $[1, 0, \cdots, 0]$.

We define the map $\pi_1 : M_{m,k} \to Y_{m,k}$ by

$$\pi_1(([Z_1, \cdots, Z_m], (\sigma, \mu))) = [\mu, b^{1/k} Z_1, \cdots, b^{1/k} Z_m], \tag{4.6}$$

where $b \in \mathbb{C}$ is defined by

$$\sigma = b((Z_1)^k, \cdots, (Z_m)^k). \tag{4.7}$$

π_1 is globally well-defined, surjective and injective on the complement of E_0.

If we identify the line bundle $\mathcal{O}(-k)$ with the open subset $\{\mu \neq 0\}$ of $M_{m,k}$ and $\mathbb{C}^m/\mathbb{Z}_k$ with the open subset $\{Z_0 \neq 0\}$ of $Y_{m,k}$ via $(z_1, \cdots, z_m) \mapsto [1, z_1, \cdots, z_m]$, then π_1 restricted to $M_{m,k}\backslash E_0$ is a biholomorphism onto $Y_{m,k}\backslash\{[1, 0, \cdots, 0]\}$ and $\pi_1(E_0) = [1, 0, \cdots, 0]$.

All of the manifolds $M_{m,k}$ admit Kähler metrics. Indeed, the cohomology classes of the line bundles $[E_0]$ and $[E_\infty]$ span $H^{1,1}(M_{m,k}; \mathbb{R})$ and every Kähler class α can be written uniquely as

$$\alpha = \frac{b}{k}[E_\infty] - \frac{a}{k}[E_0] \tag{4.8}$$

for constants a, b with $0 < a < b$. The first chern class

$$c_1(M_{m,k}) = \frac{m+k}{k}[E_\infty] - \frac{n-k}{k}[E_0]. \tag{4.9}$$

Hence if $1 \leq k \leq m-1$, then $M_{m,k}$ is a Fano manifold. He and Sun proved that any weighted projective space exists an orbifold Kähler (in fact is Kähler-Ricci soliton) metric ω_{orb} with positive bisectional curvature, see Theorem 1.2 in [20].

Let L be the $(-k)$ line bundle over $\mathbb{P}^{m-1}$, for $k \geq 1$. We give a description of the total space of L as follows. Writing $[Z_1, \cdots, Z_m]$ for the homogeneous coordinates on $\mathbb{P}^{m-1}$, we define

$$L = \{([Z_1, \cdots, Z_m], \sigma) \in \mathbb{P}^{m-1} \times \mathbb{C}^m | \sigma \text{ is in the line } \lambda \mapsto (\lambda(Z_1)^k, \cdots, \lambda(Z_m)^k)\}, \tag{4.10}$$

and let $P : L \to \mathbb{P}^{m-1}$ be the projection onto the first factor. Each fiber $P^{-1}([Z_1, \cdots, Z_m])$ is the line in $\mathbb{C}$. L can be given m complex coordinate charts

$$U_i = \{([Z_1, \cdots, Z_m], \sigma) \in L | Z_i \neq 0\} \text{ for } i = 1, \cdots, m.$$

On U_i, we have coordinates $\omega_{(i)}^j$ for $j = 1, \cdots, m$ with $j \neq i$ and $y_{(i)}$. The $\omega_{(i)}^j$ are defined by

$$\omega_{(i)}^j = Z_j/Z_i \text{ for } j \neq i,$$

and $y_{(i)}$ by

$$\sigma = \frac{y_{(i)}}{(Z_i)^k}((Z_1)^k, \cdots, (Z_m)^k).$$

On $U_i \cap U_l$ with $i \neq l$, we have

$$\omega_{(i)}^j = \tfrac{\omega_{(l)}^j}{\omega_{(l)}^i} \text{ for } j \neq i, l, \quad \omega_{(i)}^l = \tfrac{1}{\omega_{(l)}^i} \text{ and } y_{(i)} = y_{(l)}(\tfrac{Z_i}{Z_l})^k = y_{(l)}(\omega_{(l)}^i)^k.$$

Now let E be the submanifold of L defined by the zero section of L over $\mathbb{P}^{m-1}$. Denote by $[E]$ the pull-back line bundle P^*L over L, which corresponds to the hypersurface E. Writing the transition functions of $[E]$ in $U_i \cap U_l$ as $t_{il} = (Z_i/Z_l)^k = y_{(i)}/y_{(l)}$, we have a section $\tilde{s}$ over $[E]$ given by

$$s_i : \ U_i \to \mathbb{C}, \quad s_i = y_i.$$

We can define a Hermitian metric $\tilde{h}$ on the fibers of $[E]$ by

$$h_i = \frac{(\Sigma_{j=1}^m |Z_j|^2)^k}{|Z_i|^{2k}} \text{ on } U_i.$$

Namely, $\tilde{h}$ is the pull-back of h_{FS}^{-k} where h_{FS} is the Fubini-Study metric on $\mathcal{O}(1)$. We have

$$|\tilde{s}|^2_{\tilde{h}} = |y_{(i)}|^2 \frac{(\Sigma_{j=1}^m |Z_j|^2)^k}{|Z_i|^{2k}} \quad \text{on } U_i. \tag{4.11}$$

If we denote $r^2 = \sum_{i=1}^m |z^i|^2$, then we have

$$\pi_1^* r^{2k} = |\tilde{s}|^2_{\tilde{h}} \tag{4.12}$$

on L.

Let ω_e be the standard orbifold metric on $\mathbb{C}^m/\mathbb{Z}_k$, which lifts to the Euclidean metric on $\mathbb{C}^m$, we write ω_e as

$$\omega_e = \frac{\sqrt{-1}}{2\pi} \Sigma_i dz^i \wedge d\overline{z^i}. \tag{4.13}$$

Denote ω_F be the metric ω_X in Lemma 2.3 in [40], it is a Kähler metric on $M_{m,k}$. We will work in a local uniformizing chart around the orbifold point $p \in Y_{n,k}$, which we identify with the unit ball D_1 in $\mathbb{C}^m$. Then we know that ω_{orb} is uniformly equivalent to the Euclidean metric ω_e on D_1. We write D_R for the ball in $\mathbb{C}^m$ of radius $R > 0$. Then from the section 2 in [40], on $\pi_1^{-1}(D_1 \backslash \{0\})$ we have

$$k|\tilde{s}|^{2(k-1)/k}_{\tilde{h}} \pi_1^* \omega_e \leq \omega_F \leq \frac{C}{|\tilde{s}|^{2/k}_{\tilde{h}}} \pi_1^* \omega_e \tag{4.14}$$

for some uniform constant $C > 0$. Hence on $\pi_1^{-1}(D_1 \backslash \{0\})$ there exists a uniform constant C such that

$$C^{-1}|\tilde{s}|^{2(k-1)/k}_{\tilde{h}} \pi_1^* \omega_{\text{orb}} \leq \omega_F \leq \frac{C}{|\tilde{s}|^{2/k}_{\tilde{h}}} \pi_1^* \omega_{\text{orb}}. \tag{4.15}$$

Now if we denote $\tilde{\omega} = \Phi^*(\text{Pr}_2^* \pi_1^* \omega_{\text{orb}} + \text{Pr}_1^* \omega_Y)$, since the bisectional curvature of ω_{orb} is positive, using the same argument of Lemma 3.3, we obtain

Lemma 4.1. *There exist uniform constants $C > 0$ and $\alpha > 0$ such that for $\omega = \omega(t)$ a solution of the Kähler-Ricci flow,*

$$\omega(t) \leq \frac{C}{|s|^{2\alpha}_h |\tilde{s}|^{2/k}_{\tilde{h}}} \tilde{\omega}. \tag{4.16}$$

Using the same notations as in Lemma 3.4, then we have

Lemma 4.2. *For $\omega = \omega(t)$ a solution of the Kähler-Ricci flow. Then there exist uniform constants $C > 0$ and $R_0 = R_0(m,k) \in (0,1)$ such that on $D_{R_0} \backslash \{0\}$:*

$$|\tilde{V}|^2_\omega \leq C r^{2k/k+1}. \tag{4.17}$$

Locally, in $D_{R_0}\backslash\{0\}$ we have

$$|W|_g^2 \leq \frac{C}{r^{2/(k+1)}}, \tag{4.18}$$

for

$$W = \sum_{i=1}^{m}(\frac{x^i}{r}\frac{\partial}{\partial x^i} + \frac{y^i}{r}\frac{\partial}{\partial y^i})$$

the unit length radial vector field with respect to g_e, where $z^i = x^i + \sqrt{-1}y^i$.

Proof. We using (4.14) and using the same argument in the proof of of Lemma 3.4, we can obtain the Lemma. □

Using the same argument in the proof of Lemma 3.5, we can estimate on the lengths of spherical and radial paths in the punctured ball $D_{R_0}\backslash\{0\}$.

Lemma 4.3. *We have*

1. *For any $y \in Y$ and for $0 < r < R_0$, the diameter of the $2m-1$ sphere S_r of radius r in D_{R_0} centered at the origin with the metric induced from $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above, independent of r and y.*
2. *For any $z \in D_{R_0}\backslash\{0\}$, the length of a radial path $\gamma(\lambda) = \lambda z$ for $\lambda \in (0,1]$ with respect to $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above by a uniform constant multiple of $|z|^{k/(k+1)}$.*

Hence the diameter of $D_{R_0}\backslash\{0\}$ with respect to $\omega|_{\pi^{-1}(y)}$ is uniformly bounded from above and

$$\mathrm{diam}(\pi^{-1}(y), \omega|_{\pi^{-1}(y)}) \leq C.$$

Then using the same argument in Section 3, we can prove the diameter of the fiber along the metrics $g(t)$ tend to zero, then we obtain

Theorem 4.1. *Let (X, Y, π, F) be a Fano bundle with F is $M_{m,k}(1 \leq k < m)$, ω_Y be a Kähler metric on Y and ω_0 be a Kähler metric on X. Assume $\omega(t)$ is a solution of the Kähler-Ricci flow (1.1) for $t \in [0,T)$ with initial metric ω_0 and $[\omega_0] - 2\pi T c_1(X) = [\pi^*\omega_Y]$, then we have*

1. *$\mathrm{diam}(X, \omega(t)) \leq C$ for some uniform constant $C > 0$;*
2. *There exists a sequence of times $t_i \to T$ and a distance function d_Y on Y (which is uniformly equivalent to the distance induced by ω_Y, such that $(X, \omega(t_i))$ converge to (Y, d_Y) in the Gromov-Hausdorff sense.*

Acknowledgements

This is part of the first author's thesis at Rutgers University. We thank Professor Jian Song for suggesting the problem and many helpful suggestions. The work is carried out during the second author's visit at Rutgers University. He would like to thank the Department of Mathematics for its hospitality. His research is partially supported by NSFC Grant No.11301017, Research Fund for the Doctoral Program of Higher Education of China and the Fundamental Research Funds for the Central Universities and the Scholarship from China Scholarship Council.

References

[1] Aubin, T., Équations du type Monge-Ampère sur les variétés kählériennes compactes, Bull. Sci. Math., (2) 102(1978), 63-95.

[2] Birkar, C., Cascini, P., Hacon, C. and McKernan, J., Existence of minimal models for varieties of log general type, J. Amer. Math. Soc., 23(2010), 405-468.

[3] Calabi, E., Extremal Kähler metrics, Seminar on Differential Geometry, Annals of Mathematical Studies 102 (Princeton University Press, Princeton, NJ, 1982) 259-290.

[4] Cao, H., Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds, Invent. Math. 81(1985), 359-372.

[5] Chen, X. X. and Tian, G., Ricci flow on Kähler-Einstein surfaces, Invent. Math., 147(2002), 487-544.

[6] Chen, X. X. and Tian, G., Ricci flow on Kähler-Einstein manifolds, Duke Math. J. 131 (2006), 17-73.

[7] Chen, X. X. and Wang, B., The Kähler-Ricci flow on Fano manifolds (I), J. Eur. Math. Soc., 14(2012), 2001-2038.

[8] Chow, B., Chu, S.-C., Glickenstein, D., Guenther, C., Isenberg, J., Ivey, T., Knopf, D., Lu, P., Luo, F. and Ni, L., The Ricci flow: techniques and applications. Part I. Geometric aspects. Mathematical Surveys and Monographs, 135, American Mathematical Society, Province, RI, 2007.

[9] Collins, T. and Tosatti, V., Kähler currents and null loci, Invent. Math., 202(2015), 1167-1198.

[10] Feldman, M., Ilmanen, T. and Knopf, D., Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons, J. Differential Geom., 65 (2003), no. 2, 169-209.

[11] Fong, F. T.-H, Kähler-Ricci flow on projective bundles over Kähler-Einstein manifolds, Trans. Amer. Math. Soc., 366(2014), no. 2, 563-589.

[12] Fong, F. T.-H, On the collapsing rate of the Kähler-Ricci flow rate of the Kähler-Ricci flow with finite-time singularity, J. Geom. Anal., 25(2015), no. 2, 1098-1107.

[13] Fong, F. T.-H. and Zhang, Z., The collapsing rate of the Kähler-Ricci flow with regular infinite time singularity, J. Reine Angew. Math. 703 (2015), 95-113.

[14] Griffiths, P. and Harris, J., Principles of algebraic geometry, Pure Appl. Math., Wiley-Interscience, New York, 1978.

[15] Gill, M., Collapsing of products along the Kähler-Ricci flow, Trans. Amer. Math. Soc., 366 (2014), no.7, 3907-3924.

[16] Guo, B., On the Kähler-Ricci flow on projective manifolds of general type, arXiv:1501.04239.

[17] Guo, B., Song, J. and Weinkove, B., Geometric convergence of the Kähler-Ricci flow on complex surfaces of general type, arXiv:1505.00705, to appear in IMRN.

[18] Hamilton, R. S., Three manifolds with positive Ricci curvature, J. Differential Geom., 17 (1982), no.2, 255-306.

[19] Hacon, C. and McKernan, J., Existence of minimal models for varieties of log general type. II, J. Amer. Math. Soc., 23(2010), 469-490.

[20] He, W. and Sun, S., Frankel conjecture and Sasaki geometry, Adv. Math., 291 (2016), 912-960.

[21] Ilmanen, T. and Knopf, D., A lower bound for the diameter of solutions to the Ricci flow with nonzero $H^1(M^n;\mathbb{R})$, Math. Res. Lett., 10 (2003), no 2-3, 161-168.

[22] La Nave, G. and Tian, G., Soliton-type metrics and Kähler-Ricci flow on symplectic quotients, J. Reine Angew. Math. 711(2016), 139-166.

[23] Perelman, G., The entropy formula for the Ricci flow and its geometric applications, arXiv: math.DG/0211159.

[24] Perelman, G., Ricci flow with Surgery on Three-manifolds, arXiv: math.DG/0303109.

[25] Perelman, G., Finite Extinction Time for the Solutions to the Ricci flow on Certain Three-Manifolds, arXiv:math.DG/0307245.

[26] Phong, D. H. and Sturm, J., On stability and the convergence of the Kähler-Ricci flow, J. Diff. Geom., 72(2006),149-168.

[27] Phong, D. H., Song, J., Sturm, J. and Weinkove, B., The Kähler-Ricci flow and the $\overline{\partial}$-operator on vector fields, J. Diff. Geom., 81(2009), 631-647.

[28] Phong, D. H., Song, J., Sturm, J. and Weinkove, B., The Kähler-Ricci flow with positive bisectional curvature, Invent. Math., 173(2008), 651-665.

[29] Phong, D. H., Song, J., Sturm, J. and Weinkove, B., On the convergence of the modified Kähler-Ricci flow and solitons, Comment. Math. Helv. 86(2011), 91-112.

[30] Sesum, N. and Tian, G., Bounding scalar curvature and diameter along the Kähler-Ricci flow (after Perelman), J. Inst. Math. Jussieu 7 (2008), 575-587.

[31] Song, J., Finite time extinction of the Kähler-Ricci flow, Math. Res. Lett., 21(2014), no. 6, 1435-1449.

[32] Song, J., Ricci flow and birational surgery, preprint, arxiv: 1304.2607.

[33] Song, J., Székelyhidi, G. and Weinkove, B., The Kähler-Ricci flow on projective bundles, IMRN, 2013, No.2, 243-257.
[34] Song, J. and Tian, G., The Kähler-Ricci flow on surfaces of positive Kodaira dimension, Invent. Math., 170(2007), No.3, 609-653.
[35] Song, J. and Tian, G., Canonical measures and Kähler-Ricci flow, J. Amer. Math. Soc., 25(2012), No.2, 303-353.
[36] Song, J. and Tian, G., The Kähler-Ricci flow through singularities, arxiv: 0909.4898, to appear in Invent. Math.
[37] Song, J. and Tian, G., Bounding scalar curvature for global solutions of the Kähler-Ricci flow, American Journal of Mathematics, 138 (2016), no.2.
[38] Song, J. and Weinkove, B., The Kähler-Ricci flow on Hirzebruch surfaces, J. Reine Angew. Math., 659(2011), 141-168.
[39] Song, J. and Weinkove, B., Contracting exceptional divisors by the Kähler-Ricci flow, Duke Math. Jour., 162(2013), No.2, 367-415.
[40] Song, J. and Weinkove, B., Contracting exceptional divisors by the Kähler-Ricci flow II, Proc. London Math. Soc., 108(2014), No. 3, 1529-1561.
[41] Song, J. and Weinkove, B., Introduction to the Kähler-Ricci flow, Chapter 3 of 'Introduction to the Kähler-Ricci flow', eds S. Boucksom, P. Eyssidieux, V. Guedj, Lecture Notes Math. 2086, Springer 2013.
[42] Song, J. and Yuan, Y., Metric flips with Calabi ansatz, Geom. Funct. Anal., 22(2012), no. 1, 240-265.
[43] Szekelyhidi, G., The Kähler-Ricci flow and K-polystability, Amer. J. Math., 132 (2010), 1077-1090.
[44] Tian, G., New results and problems on Kähler-Ricci flow, Astérisque No. 322(2008), 71-92.
[45] Tian, G., Finite-time singularity of Kähler-Ricci flow, Discrete Contin. Dyn. Syst., 28(2010), no. 3, 1137-1150.
[46] Tian, G., Zhang, S. J., Zhang, Z. L. and Zhu, X. H., Perelman's entropy and Kähler-Ricci flow on a Fano manifold, Trans. Amer. Math. Soc. 365(2013), 6669-6695.
[47] Tian, G. and Zhang, Z., On the Kähler-Ricci flow on projective manifolds of general type, Chinese Ann. Math. Ser. B 27(2006), No. 2, 179-192.
[48] Tian, G. and Zhang, Z.-L., Convergence of Kähler-Ricci flow on lower dimensional algebraic manifolds of general type, arXiv: 1505.01038.
[49] Tian, G. and Zhang, Z.-L., Regularity of Kähler-Ricci flows on Fano manifolds, Acta. Math., 216(2016), 127-176.
[50] Tian, G. and Zhu, X. H., Convergence of Kähler-Ricci flow, J. Amer. Math. Soc., 20(2007) 675-699.
[51] Tian, G. and Zhu, X. H., Convergence of the Kähler-Ricci flow on Fano manifolds, J. Reine Angew. Math. 678(2013), 223-245.
[52] Tosatti, V., Weinkove, B. and Yang, X., The Kähler-Ricci flow, Ricci flat metrics and collapsing limits, arXiv: 1408.0161.
[53] Tosatti, V. and Zhang, Y., Infinite time singularities of the Kähler-Ricci flow, Geom. Topol., 19(2015), 2925-2948.
[54] Tosatti, V. and Zhang, Y., Finite time collapsing of the Kähler-Ricci flow on threefolds, arXiv: 1507.08397.
[55] Tsuji, H., Existence and degeneration of Kähler-Einstein metrics on minimal algebraic varieties of general type, Math. Ann., 281 (1988), no. 1, 123-133.
[56] Yau, S., On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I, Comm. Pure Appl. Math. 31 (1978), 339-411.
[57] Zhang, Z., Scalar curvature behavior for finite-time singularity of Kähler-Ricci flow, Michigan Math. J., 59(2010), no. 2, 419-433.
[58] Zhang, Z., Ricci lower bound for Kähler-Ricci flow, Commun. Contemp. Math., 16(2014), no. 2, 1350053, 11pp.

Department of Mathematics, Rutgers University, Piscataway, NJ 08854, USA
E-mail address: xf35@scarlatmail.rutgers.edu

School of Mathematics and Systems Science, Beihang University, Beijing 100191, P.R. China
E-mail address: shijinzhang@buaa.edu.cn